

## Procedure

# Manual enzymatic digestion using the dissociation enzyme mix A (lung), B (liver) or C (brain)

## Introduction

This procedure provides recommendations for performing enzymatic dissociation in single-cell workflows using manual tissue dissociation at 4°C or 37°C. For automated tissue dissociation with the Omics bundle (VIA Extractor™\* tissue disaggregator, VIA Freeze™ Uno controlled-rate freezer, and Omics clamp) and Omics pouch, please refer to the semi-automated enzymatic digestion [procedure](#).

Source	Material
<b>Provided in the dissociation enzyme mix A</b>	<ul style="list-style-type: none"><li>Enzyme 3</li><li>Enzyme 5</li><li>Enzyme 6</li></ul>
<b>Provided in the dissociation enzyme mix B</b>	<ul style="list-style-type: none"><li>Enzyme 3</li><li>Enzyme 5</li></ul>
<b>Provided in the dissociation enzyme mix C</b>	<ul style="list-style-type: none"><li>Enzyme 4</li><li>Enzyme 5</li></ul>
<b>Equipment provided by the user</b>	<ul style="list-style-type: none"><li>Micropipettes</li><li>Thermo-mixer or heating device that can heat to 37°C</li><li>Pipette controller</li></ul>
<b>Consumables provided by the user for the lung</b>	<ul style="list-style-type: none"><li>Mouse lung tissue sample: 110-150 mg cut into 5 mm pieces</li><li>50 mL conical centrifuge tubes</li><li>15 mL centrifuge tube</li><li>70 µm cell strainer</li><li>40 µm cell strainer</li><li>1000 µL wide-bore pipette tips</li><li>25 mL, 10 mL, and 5 mL serological pipettes</li><li>Red blood cell removal/lysis kit (available from a third party)</li><li>Water for cell culture</li><li>Dulbecco's phosphate-buffered saline (DPBS)</li><li>Micropipette tips</li><li>Bovine serum albumin (BSA)</li><li>Fetal bovine serum (FBS)</li><li>Sterile 25 mL reservoir</li><li>Debris removal solution (available from a third party)</li><li>0.2 µm sterile filter</li></ul>

\*For research use only.

## Required materials

The following materials are typically required in combination with the dissociation enzyme mix for tissue dissociation.

Source	Material
<b>Consumables provided by the user for the liver</b>	<ul style="list-style-type: none"><li>Mouse liver tissue sample: 400- 500 mg with gall bladder removed</li><li>50 mL sterile conical centrifuge tubes</li><li>15 mL centrifuge tube</li><li>70 µm cell strainer</li><li>1000 µL wide-bore pipette tips</li><li>25 mL, 10 mL, and 5 mL serological pipettes</li><li>Red blood cell removal/lysis kit (available from a third party)</li><li>Water for cell culture</li><li>Dulbecco's modified eagle medium (DMEM) with high glucose</li><li>Dulbecco's phosphate-buffered saline (DPBS)</li><li>Micropipette tips</li><li>Bovine serum albumin (BSA)</li><li>Fetal bovine serum (FBS)</li><li>Sterile 25 mL reservoir</li><li>Debris removal solution (available from a third party)</li><li>0.2 µm sterile filter</li></ul>
<b>Consumables provided by the user for the brain</b>	<ul style="list-style-type: none"><li>Adult mouse brain tissue samples up to 500 mg</li><li>50 mL conical centrifuge tubes</li><li>15 mL centrifuge tube</li><li>70 µm cell strainer</li><li>1000 µL wide-bore pipette tips</li><li>25 mL, 10 mL, and 5 mL serological pipettes</li><li>Red blood cell removal/lysis kit (available from a third party)</li><li>Water for cell culture</li><li>Hank's Balanced Salt Solution (HBSS)</li><li>10x Dulbecco's phosphate-buffered saline (DPBS)</li><li>Micropipette tips</li><li>Bovine serum albumin (BSA)</li><li>Fetal bovine serum (FBS)</li><li>25 mL reservoir</li><li>Cytiva's Percoll™ solution</li><li>0.2 µm sterile filter</li></ul>

## Preparation

### Note:

- For cold tissue dissociation, keep all reagents and tissue on ice, prechill the centrifuge to 4°C, and carry out benchwork on ice.
- When following the warm tissue dissociation protocol, keep all reagents and tissue at room temperature (RT) and perform all benchwork at RT.
- Tissue should be stored in the storage solution. See the next section for details on which storage solution is required for each tissue type.

1. Prepare the quench solution and the resuspension solution as detailed in the table below:

Kit-Tissue	Quench solution	Resuspension solution	Storage solution
Kit A-Lung	DPBS+10% (v/v) FBS	DPBS+0.5% (w/v) BSA	DPBS
Kit B-Liver	DMEM +10% (v/v) FBS	DPBS+0.5% (w/v) BSA	DMEM
Kit C-Brain	DPBS+10% (v/v) FBS	DPBS+0.5% (w/v) BSA	HBSS

2. Prepare the dissociation enzyme mix A, B, or C, as required:

- a. Reconstitute the enzymes according to the *Instructions for Use* for the specific enzyme mix, found in the related documents section [here](#).

- b. Prepare the enzyme mixture according to the table below:

	Mix A - lung	Mix B - liver	Mix C - brain
Reagent	Volume for 1 sample	Volume for 1 sample	Volume for 1 sample
Enzyme 3	2 mL	1 mL	-
Enzyme 4			2 mL
Enzyme 5	20 µL	10 µL	20 µL
Enzyme 6	300 µL	-	-
DMEM	-	2990 µL	-
Total volume	2.32 mL	4 mL	2.02 mL

### Note:

- To calculate volumes for multiple samples, multiply the volume for one sample by the number of samples and add 3%.
- For cold tissue dissociation, keep enzyme mixes on ice. When following the warm tissue dissociation protocol, warm the enzyme mix at 37°C for 30 minutes.
- Recommended sample sizes, buffers, dissociation temperatures, and time for manual dissociation are provided in the table below:

Mix and tissue type	Size/weight	Storage solution	Quench solution	Resuspension buffer	Cell strainer size	Temperature (°C)	Time
Mix A - lung	Mouse lungs, 110 to 150 mg	DPBS	DPBS + 10% FBS	DPBS + 0.5% BSA	70 µm and 40 µm	37	30 min
Mix B - liver	Liver tissue, 400 to 500 mg	DMEM	DMEM + 10% FBS	DPBS + 0.5% BSA	70 µm	37	30 min
Mix C - brain	1 adult brain up to 500 mg	HBSS	DPBS + 10% FBS	DPBS + 0.5% BSA	70 µm	4 and 37	30 min

## Manual dissociation protocol

1. Mince the tissue by cutting finely with a scalpel.
2. Transfer the minced tissue to a 15 mL centrifuge tube.
3. Add the enzyme mix to the tissue.
4. Triturate the sample 10 times with a wide-bore 1 mL pipette tip (T=0, 10, 20, and 30 min).
5. Incubate for 30 minutes, repeating step 4 every 10 minutes.
  - For warm dissociation: Incubate on a dry bath/block incubator at 37°C.
  - For cold dissociation: Incubate on ice.
6. Wet the 70 µm cell strainer with 2 mL of quench solution.
7. Pass the digested material through the cell strainer into a 50 mL conical centrifuge tube.
8. Rinse the digestion vessel with 2 mL of quench solution and pass the resultant solution over the strainer.
9. Use the plunger from a 5 mL syringe (or similar) to gently push the tissue material remaining in the filter through the filter.
10. Wash filter 3 times, with 1 mL quench solution.
11. Centrifuge the samples at 300 x g for 10 minutes.
12. Remove the supernatant and resuspend the pellet in resuspension solution.
13. For optimal results, additional steps, including red blood cell (RBC) removal or debris removal might be required. For brain post processing, refer to 'Myelin removal for brain tissue'.

## Post-processing for lung tissue

1. For dissociation of lung tissue, the following steps are optional to remove cell clumps. After RBC removal, add 1 mL of resuspension solution to the cell pellet at RT.
2. Mix thoroughly by pipetting.
3. Place a 40 µm strainer on separate 50 mL centrifuge tubes.
4. Pass the sample over the 40 µm cell strainer and wait until the sample has passed through.

## Post-processing for liver tissue

For optimal results, it is recommended to perform RBC removal followed by debris removal.

## Myelin removal for brain tissue

1. Perform RBC removal as per the manufacturer's protocol
2. Dilute 2 mL of 10x DPBS in 18 mL of Percoll solution. Keep the mixture on ice.
3. Add 15 mL of the solution from step 2 to 35 mL of ice-cold 1x DPBS to make 27% Percoll solution. Keep on ice.
4. Resuspend the cell pellet after RBC removal in resuspension buffer. Pipette mix gently with a 1 mL wide-bore tip.
5. Measure dissociated brain cell suspension using a 2 mL serological pipette, note the volume of the cell pellet, and transfer the pellet to a 15 mL centrifuge tube.
6. Add 6 mL of 27% Percoll solution per 1 mL of brain sample. Mix gently by inverting the tube.
7. Centrifuge at 4°C at 700 x g for 10 minutes with the brake off.
8. Carefully remove and discard the upper myelin layer using a 2 mL serological pipette.
9. Remove and discard the remaining supernatant using a serological pipette.
10. Resuspend cells in 5 mL DPBS + 0.5% BSA and slowly pipette the pellet to wash.
11. Change centrifuge deceleration settings to maximum and centrifuge at 300 x g for 10 minutes.
12. Remove the supernatant and resuspend the pellet in 1 mL DPBS + 0.5% BSA.



Refer a colleague to claim your full FREE kit or request a sample pack to get started. Scan to learn more or click here! [cytiva.link/gz1v1](https://cytiva.link/gz1v1)

**cytiva.com**

Cytiva and the Drop logo are trademarks of Life Sciences IP Holdings Corp. or an affiliate doing business as Cytiva. Percoll, VIA Extractor, and VIA Freeze are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva.

Any other trademarks are the property of their respective owners.

© 2026 Cytiva

For local office contact information, visit [cytiva.com/contact](https://cytiva.com/contact)

CY54301-06Feb26-CP

 **cytiva**