

IgG Sepharose 6 Fast Flow

Instructions for Use

IgG Sepharose™ 6 Fast Flow, from Cytiva, is based on the rigid Sepharose 6 Fast Flow matrix, with human IgG covalently coupled to it. The improved mechanical characteristics of this fast flow medium allows high flow rates to be used for rapid and convenient single step purification of protein A fusion protein conjugates produced in prokaryotic expression systems. IgG Sepharose can be used in combination with Calmodulin Sepharose for tandem affinity chromatography (TAP) purification of protein complexes. Characteristics of IgG Sepharose 6 Fast Flow are listed in the table below.

Table 1. Characteristics of IgG Sepharose 6 Fast Flow

Ligand:	human IgG
Binding capacity:	> 2 mg protein A/ml medium at pH 7.5
Mean particle size:	90 µm
Bead structure:	highly cross-linked 6% agarose
Maximum flow rate ¹ :	300 cm/h (98 ml/min), using XK 50/30 column with 15 cm bed height, run at room temperature with aqueous buffer
Recommended flow rates ¹ :	Sample application: < 150 cm/h (49 ml/min using XK 50/30 column)
Storage temperature:	2°C to 8°C
Storage buffer:	50 mM K ₂ PO ₄ , 1 mM MgCl ₂ pH 7.2 in 20% ethanol or 50 mM Tris buffer, 150 mM NaCl, 0.05% Tween™ 20, pH 7.6 (TST) in 20% ethanol

¹ H₂O at room temperature

Protocol

Buffers

- Tris-saline Tween 20 (TST): 50 mM Tris buffer, pH 7.6, 150 mM NaCl and 0.05% Tween 20.
- 0.5 M CH_3COOH (HAc) adjusted to pH 3.4 with $\text{CH}_3\text{COONH}_4$ (NH_4Ac).
- 5 mM NH_4Ac , pH 5.0.

Procedure



WARNING

When using hazardous chemicals, take all suitable protective measures, such as wearing protective glasses and gloves resistant to the chemicals used. Follow local regulations and instructions for safe operation and maintenance of the system.

Column Packing and Washing

Step	Action
------	--------

- | | |
|---|--|
| 1 | IgG Sepharose 6 Fast Flow is supplied as a suspension in 50mM K_2PO_4 , pH 7.2, 1mM MgCl_2 , 20% ethanol as preservative. |
| 2 | Pack the IgG Sepharose 6 Fast Flow suspension in any Cytiva column of suitable size. The column may be operated by gravity feed. |

Step	Action
------	--------

- | | |
|---|--|
| 3 | Wash the medium with at least 5 bed volumes TST prior to use in order to remove any traces of ethanol. |
|---|--|

Equilibration and Sample Application

Step	Action
------	--------

- | | |
|---|--|
| 1 | Equilibrate the column with 2–3 bed volumes each of <ul style="list-style-type: none">• 0.5 M HAc, pH 3.4• TST• 0.5 M HAc, pH 3.4• TST |
| 2 | Check pH of the eluate with pH paper (should be neutral) and adjust pH of the sample, cell supernatant or clarified growth medium, if necessary. Apply the sample to the column. |

Washing and Elution

Step	Action
------	--------

- | | |
|---|--|
| 1 | Wash the medium with <ul style="list-style-type: none">• 10 bed volumes TST• 2 bed volumes 5mM NH₄Ac, pH 5.0 |
| 2 | Elute the sample with 0.5 M HAc, pH 3.4. |

Step	Action
3	Collect aliquots in polypropylene microcentrifuge tubes ($A_{280}=1.0$ for 2.6 mg protein/ml).
4	The samples can be lyophilized directly without prior dialysis.
5	Analyze samples by gradient sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE).
Note: <i>This method gives a concentrated eluate and can only be used if the fusion product is stable under these conditions. For small scale purifications, gravity flow suffices throughout the procedure.</i>	

Alternative Elution Buffers

- 0.1 M glycine-HCl, pH 3.0.
- 0.5 M lithium diiodosalicylate dissolved in water (enzymatic activity is better preserved but some of the sample will bind to the column irreversibly). Other chaotropic salts may also be used.

Re-equilibration and Storage

- Re-equilibrate IgG Sepharose 6 Fast Flow with TST until pH of the effluent is around 7.0. This is important since IgG might denature if the chromatography medium is left standing at a low pH.

- If the medium is not going to be used for a longer period of time, wash the matrix with 5 bed volumes 20% ethanol in TST and store at 2°C to 8°C. IgG Sepharose 6 Fast Flow must not be frozen.

Shelf Life – General Guidelines

- Human IgG is covalently coupled to the Sepharose matrix via cyanogen bromide activation. Minute amounts of IgG that leak from the medium will be washed away in the initial washing step. No IgG is visible on a silver stained PhastGel™ Gradient 10–15 run on PhastSystem™. The sensitivity limit of this silver staining technique is 0.5 ng protein per band for PhastGel SDS-PAGE.
- Careful sample preparation will prolong the life of IgG Sepharose 6 Fast Flow. Avoid reducing agents since the disulphide bonds in IgG will be affected.
- Coloured samples may leave traces of pigment on the medium. This colouration does not influence the performance of IgG Sepharose 6 Fast Flow.
- The column may be operated from 4°C to room temperature, depending on the sensitivity of the protein conjugate.

Ordering Information

Product	Quantity	Product Code.
IgG Sepharose 6 Fast Flow	10 ml	17096901

Related products	Quantity	Product Code.
Calmodulin Sepharose 4B	10 ml	17052901

Literature	Quantity	Product Code.
Affinity Chromatography Handbook, Principles and Methods	1	18102229
Affinity Chromatography Columns and Media Selection guide	1	18112186

Page intentionally left blank



cytiva.com

Cytiva and the Drop logo are trademarks of Global Life Sciences IP Holdco LLC or an affiliate.

PhastGel, PhastSystem, and Sepharose are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva.

All other third-party trademarks are the property of their respective owners.

Tween is a trademark of Croda Group of Companies

© 2020 Cytiva

All goods and services are sold subject to the terms and conditions of sale of the supplying company operating within the Cytiva business. A copy of those terms and conditions is available on request. Contact your local Cytiva representative for the most current information.

For local office contact information, visit [cytiva.com/contact](https://www.cytiva.com/contact)

52208300 AK V:6 07/2020